Index to Volume 86 of THE BRYOLOGIST[1,2]

Abietinella abietina, 54
Acarospora fuscata, 16; scabrida, 16
acclimation, in lichens, 31+
acetic-haematoxylin, for cytological studies of bryophytes, 141
acid phosphatase, 129+
Actinostoma, *194*
Agyrophora lyngei, 16
Alectoria ochroleuca, 16, 40
alkaline phosphatase, 129+
ALLEN, B. H. Isothecium myosuroides Brid. and I. stoloniferum Brid.: A Quanti..ative Study, 358
ALLEN, B. H. & J. A. SNIDER. Additions to the Nicaraguan Moss Flora, 378
Amblyodon dealbatus, 54
Amblystegiella jungermannioides, *60*
Amblystegium cf. serpens, 372; compactum, *60*; juratzkanum, *60*, 127; polygamum, 336; saxatile, *60*; serpens, 54, 110; stellatum, 336; tibetanum, 127; varium, 4, 127, 278
Amphidium lapponicum, 54, 334; mougeotii, 54
Anaptychia palmulata, 118
Anastrophyllum minutum, 337
ANDERSON, LEWIS E. Charles M. Boardman (1903–1983), 162
ANDO, H. & M. HIGUCHI. Homomallium sharpii, a New Moss Species from Mexico, 374
Andreaea blyttii, 54; crassinervia, 54; nivalis, 55; obovata, 333; papillosa, 333; rothii, 55; rupestris, 55, 333
ANDRUS, R. E. Two New North American Sphagnum Species in Section Subsecunda, 257
Anisothecium grevilleanum, *60*; palustre, *60*; schreberianum, *60*
Anoectangium bellii, 110
Anomobryum concinnatum, *60*; filiforme, 55
Anomodon attenuatus, 55, 279; rostratus, 55
Anthelia juratzkana, 334+
Antitrichia curtipendula, 55
Aongstroemia longipes, 55
ARCHER, A. W. Cladonia chlorophaea sens. lat. in Southeastern Australia, 251
Archidium ohioense, 270
Arctoa anderssonii, 337+; fulvella, 55; hyperborea, 339
Arctomia delicatula, 17; interfixa, 17
Arkansas, new moss records, 357+
Asahinea chrysantha, 16+; scholanderi, 16+
Ascidiota, 144
Aspicilia arctica, comb. nov., 14+; **bennettii**, comb. nov., 14+; cinerea, 17; myrini, 17; nikrapensis, 17; rosulata, 17; stygioplaca, 17
Asterella lindenbergiana,115
Astomum muhlenbergianum, 4
Athalamia hyalina, 115
atranorin, 21, 64+, 251
Atrichum altecristatum, 55; oerstedianum, 55; tenellum, 55; undulatum, 1+, 55, 333+
Aulacomnium androgynum, 55; palustre, 55, 260, 333; turgidum, 55, 333
Australia, new lichen records, 251+, 347+; southeastern, Cladonia chlorophaea sens. lat. in, 251

Baeomyces carneus, 17; placophyllus, 17
Barbilophozia binsteadii, 333+; floerkei, 339; hatcheri, 339; lycopodioides, 334+
Barbula asperifolia, *60*; brevifolia, 336; constricta, *271+*; constricta var. flexicuspis, *272*; convoluta, 4, 55; fallax, 55; flexicuspes, *271+*; fusca, *271+*; icmadophila, 55; orizabensis, 156; reflexa, 55; rigidula, *60*; sprengelii, *134*; stenotheca, *156*; subcontorta, *271+*; unguiculata, 4, 55
BARTLETT, J. K., see HORTON, D. G. & J. K. BARTLETT
BARTLETT, J. K., see VITT, D. H. & J. K. BARTLETT
Bartramia breviseta, 337; ithyphylla, 55, 333+; pomiformis, 55, 114
BAUER, C. R., see CROSBY, M. R. & C. R. BAUER
BAUER, C. R. & M. R. CROSBY. Recent Literature on Mosses—126–127: 300, 393
Bellincinia, 144
BISCHLER, H. Charles Hébant (Mai 1941–1982), 158
Blasia pusilla, 241
blepharoplast structure, in Sphaerocarpos and Riella, 97
Blepharostoma trichophyllum, 333+
Blindia acuta, 55, 334; calcarea, 111; magellanica, 110
Boardman, Charles M. necrology, 162
bourgeanic acid, 64+, 254
BOWERS, F. D., see MARAVOLO, N. C., F. D. BOWERS & S. FRECKMANN
Brachymenium preissianum, 110; stystilum, 378
Brachythecium albicans, 55; campestre, 55; collinum, 114; curtum, 55; erythrorrhizon, 55, 337; glaciale, 334+; oxycladon, 55; plumo-

---

[1] Compiled by P. M. Eckel, Clinton Herbarium, Buffalo Museum of Science, Buffalo, NY 14211.

[2] New scientific names and new combinations are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign; italicized page numbers indicate taxa in synonymy.

sum, 55; populeum, 55; reflexum, 55; rivulare, 55; rutabulum, 55, 278; salebrosum, 4, 55; starkei, 55; turgidum, 55; velutinum, 55
BRASSARD, G. R. Checklist of the Mosses of the Island of Newfoundland, Canada, 54
BRASSARD, G. R. (review) R. R. Ireland. Moss Flora of the Maritime Provinces, 404
BRASSARD, G. R. & T. HEDDERSON. The Distribution of Mielichhoferia macrocarpa, a North American Endemic Moss, 273
Brotherella delicatula, *60*; recurvans, 55
BROWN, R. C., see CAROTHERS, Z. B., R. C. BROWN & J. G. DUCKETT
BROWN, R. C., Z. B. CAROTHERS & J. G. DUCKETT. Comparative Spermatogenesis in the Sphaerocarpales. II. Blepharoplast Structure in Geothallus tuberosus Campb., 234
BRUGGEMAN-NANNENGA, M. A. (review) R. E. Magill. Flora of Southern Africa. Bryophyta. Part 1, Mosses. Fascicle 1, Sphagnaceae-Grimmiaceae, 403
Bryhinia novae-angliae, 55
Bryoerythrophyllum ferruginascens, 271; jamesonii, 138; recurvirostrum, 334+, 372; recurvum, 271; rubrum var. minus, *271*; tenerrimum, *271*
Bryoria chalybeiformis, 17; nitidula, 17
Bryum aeneum, 55; algovicum, 4, 55, 268; alpinum, 55; angustirete, *60*; antarcticum, 2; apiculatum, 270; archangelicum, 268; arcticum, 55, 268, 336; argenteum, *2*, 55, 335; billardieri, 378; bimum, 55; caespiticium, 4, 55; calophyllum, 55; capillare, 55; cernuum, *60*; cirrhatum, 60; creberrimum, 1+, 55+; gemmiparum, 55; harriottii, 110; inclinatum, *60*, 333; jan-mayense, 332+; klinggraeffi, 4; knowltonii, 55; lacustre, 333; lapponicum, 55; lepidum, 268; marratii, 55; micro-erythrocarpum, 110, 357; miniatum, 55; mamillatum, 268; muehlenbeckii, 55; neodamense, 278; oelandicum, 268; ovatum, 55; pallens, 55, 268, 337; pallescens, 55, 268; pendulum, *268*; pseudotriquetrum, 4, 55, 334; purpurascens, 268; oxystegium, 268; reedei, 268; salinum, 55, 336; sandbergii, 2; stenocarpum, 268; stenotrichum, 55; subnitidulum, 332; tortifolium, 55; uliginosum, 55; warneum, 266+, 267*; weigelii, 55; wrightii, 55
Buellia aethalea, 17; aethaleoides, 18; alboatra, 18; baltica, 18; immersa, 17; malmei, 18; minutissima, 18; papillata, 18; sororioides, 17
Buxbaumia aphylla, 55, 124, 372; himalayensis, 121; indusiata, 124; minakatae, 55; piperi, 121; viridis, 60

Callicladium haldanianum, 55
Calliergon cordifolium, 55; giganteum, 55; richardsonii, 55; sarmentosum, 55, 333+; stramineum, 5, 333; trifarium, 55; wickesiae, 55
Calliergonella cuspidata, 55
Caloplaca arizonica, 249+; cinnamomea, 18; elegans, 33, 249+; festiva, 18; holocarpa, 18; pinicola, 16+; saxicola, 118; stillicidiorum, 18; tetraspora, 18; tiroliensis, 18
Calymperes afzelii, 29; bartramii, 29; donnellii, *29*; erosum, 29; flexifolius, *30*; guildingii, 29; levyanum, 29; lonchophyllum, 29; mitrafugax, 29; nashii, *29*; nicaraguense, 29; palisotii subsp. richardii, 29; platyloma, 29; richardii, *30*; rubiginosum, 24+; rufum, *30*; smithii, 24+; tenerum var. edamense, 24+; uleanum, 29; venezuelanum, 29
Calypogeja integristipula, 115; sphagnicola, 115
Campylium chrysophyllum, 4, 55, 339; halleri, 55; hispidulum, 4, 55; polygamum, 55, 261, 336+; polygamum var. minus, 339; radicale, 55; stellatum, 55, 339
Campylopus atrovirens, 55; schimperi, 55; subulatus, 60; tallulensis, 378
Canada, checklist of the mosses of Newfoundland (island), 54+; lichens from Bathurst Inlet Region, Northwest Territories, 14+
Candelaria concolor, 118, 246+
Candelariella terrigena, 18; vitellina, 18
caperatic acid, 254, 365+
carbonic anhydrase, 129+
CAROTHERS, Z. B., see BROWN, R. C., Z. B. CAROTHERS & J. G. DUCKETT
CAROTHERS, Z. B., R. C. BROWN & J. G. DUCKETT. Comparative Spermatogenesis in the Sphaerocarpales. I. Blepharoplast Structure in Sphaerocarpos and Riella, 97
catalase, 129+
Catoscopium nigritum, 55
Cephalozia bicuspidata, 114; bifida, 336; pleniceps, 333
Cephaloziella rubella, 336; rubella var. bifida, 336
Ceratodon purpureus, 4, 55, 333+
Cetraria arenaria, isozyme banding in 129+; cucullata, 18; delisei, 18; hepatizon, 18; islandica, 18; islandica ssp. crispiformis, 18; islandica ssp. reticulata, 18; laevigata, 16+; nigricans, 18; nivalis, 18, 40; oakesiana, 365+; pinastri, 18, 365+; sepincola, 18; tilesii, 18
China, revision of Entodon species in, 193+
chromosome number, Amblystegium tibetanum, 121+; Racomitrium canescens, 121+; Stereophyllum fulvum, 121+; S. tavoyense, 121+; Buxbaumia himalayensis, 121+; Megalostylium brevisetum, 121+; Syrrhopodon gardneri, 121+; Cinclidium stygium, 55
Cinclidium subrotundum, 55
Cirriphyllum piliferum, 55
Cladina stellaris, 309; subtenuis, 118+
Cladonia alaskana, 18; amaurocraea, 18; apodocarpa, 118+; asahinae, 66+; bacillaris, 118; bellidiflora, 18; brevis, 118; caespiticia, 118; capitata, 118; cariosa, 118; carneola, 18; chlorophaea, 18, 64+, 118; chlorophaea sens. lat., 251+; clavulifera, 118; coccifera, 18; coni-

ocraea, 118; conista, 66+; conoidea, 251+; cristatella, 118; cryptochlorophaea, 251+; cyathomorpha, 66+; cylindria, 118+; decorticata, 118+; didyma, 118; ecmocyna, 18; fimbriata, 18, 64+; furcata, 118; gracilis, 18; gracilis ssp. gracilis, 18; gracilis ssp. turbinata, 18; grayi, 118, 251+; homosekikaica, 251+; macilenta, 118; macrophylla, 18; merochlorophaea, 251+, 66+; merochlorophaea var. merochlorophaea, 66+; merochlorophaea var. novochlorophaea, 251; mitis, 18, 335; parasitica, 118; phyllophora, 19; piedmontensis, 118+; pityrea, 118; pocillum, 19; polycarpoides, 118; pyxidata, 118; rangiferina, 19, 41; rei, 118+; squamosa, 118; stellaris, 19; stricta, 19; sulphurina, 19; symphycarpa, 118; uncialis, 19, 333; verticillata, 19, 118
Cladopodiella francisci, 337
Claopodium bolanderi, 114
Climacium dendroides, 56, 372
Collema fragrans, 118; furfuraceum, 41, 249+
Conardia compacta, 56
connorstictic acid, 254
Conostomum tetragonum, 56
conpsoromic acid, 254
CONRAN, J. G. & R. W. ROGERS. Lichen Succession on Leaves of Wilkiea macrophylla in Southeast Queensland, 347
constictic acid, 254
Cornicularia divergens, 19
Coscinodon cribrosus, 56
CRANDALL-STOTLER, B., see STOTLER, R. E. & B. CRANDALL-STOTLER
Cratoneuron commutatum, 60; falcatum, 56; filicinum, 56, 110, 279; filicinum var. curvicaule, 372; filicinum var. filicinum, 372
Cratoneuropsis relaxa, 110, 372
Cribrodontium, 194
CROSBY, M. R. & C. R. BAUER. Recent Literature on Mosses—124–125: 88, 176
CROSBY, M. R., see BAUER, C. R. & M. R. CROSBY
CRUM, H. (review) Sir H. Godwin. The Archives of the Peat Bogs, 402
cryptochlorophaeic acid, 251
Ctenidium molluscum, 56
CULBERSON, C. F., see CULBERSON, W. L., C. F. CULBERSON & S. A. FISCUS
CULBERSON, W. L., C. F. CULBERSON & S. A. FISCUS. A New Lecanoric Acid-Producing Usnea from Mexico, 254
Cuspidaria, *194*
Cylindrothecium, *194*; concinnum, *211*; montagnei *211*
Cymbifoliella, *194*
Cynodontium jenneri, 56, 115; polycarpon, 60; strumiferum, 56; strumulosum, 56; tenellum, 56
Cyrtomnium hymenophylloides, 56
cytology, acetic-haematoxylin method for bryophytes, 141
Dactylina arctica, 19; madreporiformis, 19; ramulosa, 19
DAMSHOLT, K. The Gemmae of Tritomaria (Saccobasis) polita (Nees) Schiffn. ssp. polita (Hepaticae), 139
DELGADILLO M., C. (review) O. de Bolòs, J. M. Montserrat, A. M. Romo & E. Serra (Eds.) IV. Simposi de Botànica Criptogàmica: Pteridophyta, Bryophyta et Lichenes, 405
Deploschistes scruposus, 19
Dermatocarpon hepaticum, 19; intestiniforme, 19; miniatum, 118
Desmatodon cernuus, 56; garberi, *134*; heimii, 56; latifolius var. latifolius, 56; latifolius var. muticus, 56; laurei, 56; leucostoma, 56; obtusifolius, 4, 279; spathulifolius, *134*; sprengelii, *134*; stomatodontus, *138*; systylius, 56; viridipilus, 110
DEWEY, R. M. Bryophytes of the Tenmile Creek Drainage, Idaho, 114
Dichelyma capillaceum, 56; falcatum, 56; pallescens, 56
Dichodontium olympicum, 114; pellucidum, 56, 334+
Dicranella cerviculata, 56; grevilleana, 56; heteromalla, 56; palustris, 56; rufescens, 115; schreberiana, 56; subulata, 56; varia, 56
Dicranodontium denudatum, 56
Dicranoweisia crispula, 56
Dicranum acutifolium, 56; angustum, 56, 333+; bonjeanii, 56; congestum, 334; congestum var. spadiceum, 336; drummondii, 60; elongatum, 56; fulvum, 56; fuscescens, 56, 115, 333+; groenlandicum, 56; leioneuron, 56; majus, 56, 333+; muehlenbeckii, 56, 336+; ontariense, 56; pallidisetum, 115; polysetum, 56; scoparium, 56, 334+; spadiceum, 56, 335+; spurium,56; sulcatum, 115; tauricum, 114; undulatum, 56; viride, 56
Didymodon asperifolius, 56; constrictus, *271*; constrictus var. flexicuspis, *272*; fallax, 4; rigidulus, 56, 279; tenerrimus, *271*; tophaceus, 336; vinealis, 271+; vinealis var. flaccidus, *271*+; vinealis var. vinealis, *272*
diffractaic acid, 254
Dimelaena oriena, 18+
Diphyscium foliosum, 56, 124
Diplophyllum taxifolium, 334+
Diploschistes scruposus, 19
Dirinaria frostii, 118
Discelium nudum, 60
Distichium capillaceum, 56, 334+; hagenii, 60; inclinatum, 56, 115
Ditrichum cylindricum, *60*, 333+; flexicaule, 56, 339; giganteum, *60*; heteromallum, 333+; lineare, 56; pusillum, 56
Dorcadion arcticum, 337
Drepanocladus aduncus, 56; badius, 56, 337; capillifolius, 44+; crassicostatus, 44+; crassicostatus, distribution map, 50; **crassicostatus**, sp.

nov., 45+; exannulatus, 56, 261, 337; exannulatus var. purpurascens, 44+, 372; exannulatus var. rotae, 44+; fluitans, 45, 56; procerus, 56; revolvens, 56, 334, 372; trichophyllus, 44+; uncinatus, 56, 333+; vernicosus, 56
Dryptodon patens, 56
DUCKETT, J. C., see CAROTHERS, Z. B., R. C. BROWN & J. G. DUCKETT
DUCKETT, J. G., see BROWN, R. C., Z. B. CAROTHERS & J. G. DUCKETT

EGAN, R. S. Recent Literature on Lichens—115–118: 81, 169, 293, 387
Encalypta alpina, 334+; ciliata, 56, 114; longicolla, 56; procera, 56, 279, 334+; rhaptocarpa, 56, 110, 114, 372; streptocarpa, 56; vulgaris, 372
Entodon **acutifolius**, 193+, 199*; aeruginosus, 211+, 227; amblyophyllus, *213*; angustifolius, *196*; attenuatus, *223*; caliginosus, 211+, 212*; challengeri, *214*; cladorrhizans, 210+, 221*; cochleatus, *232*; compressus, 211+; compressus var. compressus, 214, 215*; compressus var. **zikaiwiensis**, 214, 215*; conchophyllus, 211+, 224*; concinnus, 56, 211+, 212*; delavayi, *196*; divergens, 196+, 205*; dolichocucullatus, 196+, 199*; drummondii, *196*; eurhynchioides, *203*; excavatus, *196*; fissidens, *195*; flavescens, 211+, 228*; giraldii, 211+, 218*; griffithii, *229*; kungshanensis, 196+, 201*; latifolius, *225*; longicostatus, *223*; longifolius, 195+, 209*; luridus, 211+ 231*; macropodus, 196+, 197*; macrosporus, *223*; mairei, *225*; micropodus, 211+, 224*; microthecius, *214*; montagnei, *211*; morrisonensis, 211+, 220*; nanocarpus, *214*; obtusatus, 196+, *203*, 204*; okamurae, *232*; orthocarpus, *211*; plicatus, 196+, 207*; prorepens, 211+, 226*; pseudo-orthocarpus, *213*; purus, *225*; pylaisioides, 196+, 207*; ramulosus, *229*; rostrifolius, *229*; rubicundus, *229*; rubissimus, *229*; scabridens, 200; scariosus, 196+, 209*; schensianus, 211+, 230*; schwaegrichenii, *229*; serpentinus, *213*; smaragdinus, 211+, 217*; stenopyxis, *225*; subramulosus, *213*; sullivantii var. sullivantii, 211+, 222*; sullivantii var. versicolor, 211+, 222*; taiwanensis, 196+, 202*; variegatus, *229*; virens, 195; viridulus, 196+, 202*; zikaiwiensis, *214*; revision of Chinese species of, 193+; subg. Entodon, 195; subg. Erythropus, 195
Ephemerum spinulosum, 4
esterase, 129+
Eucladium irroratum, 110; verticillatum, 60
Eurhynchium diversifolium, 339; praelongum, 113; pulchellum, 4, 56, 339; riparioides, 56
evernic acid, 254

FAHSELT, D. & C. HAGEMAN. Isozyme Banding Patterns in Two Stands of Cetraria arenaria Kärnef., 129
FARRAR, D. R., see VAN DER LINDEN, J. O. & D. R. FARRAR
FISCUS, S. A., see CULBERSON, W. L., C. F. CULBERSON & S. A. FISCUS
Fissidens adiantoides, 56; arcticus, 334; bryoides, 4, 334; cristatus, 56; glaucullus, 270; osmundoides, 56, 334; viridulus, 56
Fistulariella almquistii, 16+
Fontinalis antipyretica, 56; dalecarlica, 56; flaccida, 56; gigantea, 56; novae-angliae, 56
fossils, bryophyte, 44+
Fossombronia incurva, 241
FRECKMANN, S., see MARAVOLO, N. C., F. D. BOWERS & S. FRECKMANN
FRISVOLL, A. A. Bryophytes from Jan Mayen, Including Twenty-five Species New to the Island, 332
Frullania asagrayana, 279; eboracensis, 279
fumarprotocetraric acid, 64+, 251+, 254
Funaria arctica, 334; flavicans, 4; hygrometrica, 56, 334; microstoma, 56
Fungaria glabra, 372

galbinic acid, 254
gemmae, of Tritomaria polita spp. acuta, 139+; of Tritomaria polita spp. polita, 139+
Geocalyx graveolens, 279
Geothallus tuberosus, blepharoplast structure in, 234+
Geothallus, status in Sphaerocarpales, 97
GLIME, JANICE M. (review) L. Vogelenzang (Ed.) Guide to the Prices of Antiquarian and Secondhand Botanical Books (1970–1979). Cryptogamic Literature (Algae, Desmids, Ferns, Fungi, Lichens, Mosses, Phytopathology, Phytoplankton), 190
Glossadelphus pernitens, *232*
glutamate dehydrogenase, 129+
grayanic acid, 251
Grimmia alpicola, 278, 338; apocarpa, 333+; calyptrata, 114; conferta, 333; diminuta, *111*; hartmanii var. anomala, 56; incurva, 56; janmayensis, 332+; olneyi, 56; ovalis, 56; tenerrima, 56; torquata, 56; trichophylla, 57; unicolor, 57
Groutiella apiculata, 378
Gyalectidium filicinum, 348+
Gymnomitrion concinnatum, 333
Gymnostomum aeruginosum, 57; calcareum, 110; recurvirostrum, 57, 279
Gyroweisia tenuis, 57

Haematomma lapponicum, 19
HAGEMAN, C., see FAHSELT, D. & C. HAGEMAN
Haplohymenium cf. pseudotriste, 372
Haplomitrium hookeri, 241
Hébant, Charles. Bibliography and necrology, 159+

HEDDERSON, T., see BRASSARD, G. R. & T. HEDDERSON
Hedwigia ciliata, 57
Helodium blandowii, 57
HENNINGS, C. J. The Cladonia chlorophaea-C. fimbriata Complex in Western Washington, 64
Herzogiella striatella, 57; turfacea, 57
Heterocladium dimorphum, 57, 334; procurrens 114
Heterodermia speciosa, 118
HIGUCHI, M., see ANDO, H. & M. HIGUCHI
HOCKENHULL, Y., see LEPP, N. W. & Y. HOCKENHULL
Homalia jamesii, *60*; trichomannoides, 57, 279
Homalothecium sericeum, 57
Homomallium loriforme, 376; mexicanum, 374+; **sharpii**, sp. nov., 374+, 375*; simlaense, 376
homosekikaic acid, 251
HONG, WON SHIC. The Genus Porella in North America West of the Hundredth Meridian, 143
HORTON, D. G. & J. K. BARTLETT. The Genus Timmia Hedwig: A Bipolar Disjunct, 370
HORTON, D. G. (review) Mitten, W. 1869. Musci Austro-Americani, 307
Huilia flavocaerulescens, 19; macrocarpa, 19; melinodes, 19
Hygroamblystegium chilense, 372; fluviatile, 57; tenax, 57
Hygrohypnum alpestre, 57; dilatatum, 57; eugyrium, 57; luridum, 57; molle, 114; montanum, 57; ochraceum, 57; polare, 334; smithii, 57; subeugyrium, 57
Hylocomiastrum pyrenaicum, *60*; umbratum, *60*
Hylocomium brevirostre, 57; pyrenaicum, 57; splendens, 57, 110+, 336; umbratum, 57
Hymenostylium recurvirostrum, 138
Hyophila, 156; compared with Neohyophila, 134+; fragilis, *134*; involuta 134+, 278; lingulata, *134*; stomatodonta, *138*; subangustifolia, *156*
Hyophilopsis lingulata, *134*; spathulifolia, *134*; stomatodonta, *138*; sprengelii, *134*
Hypnum bambergeri, 57; bridelianum, *60*; concinnum, *211*; cupressiforme var. cupressiforme, 57; cupressiforme var. filiforme, 57; cupressiforme var. lacunosum, 57; curvifolium, 57; dieckii, 115; fertile, 57; imponens, 57; lindbergii, 57, 278; orthocarpus, *211*; pallescens, 57; plicatulum, 57; pratense, 57; procerrimum, 57; protuberans, *60*; recurvatum, 57; revolutum, 57, 114; schreberi var. orthocarpus, *211*; vaucheri, 57
Hypogymnia austerodes, 19; oroarctica, 19; physodes, 19; subobscura, 19

Idaho, bryophytes of the Tenmile Creek Drainage, 114+; new state records, 114+
Iowa, ecological study of a natural prairie in northwestern, 1; lichens of eastern, 116+; new lichen records, 116+; new bryophyte records, 4
Isopterygiopsis muelleriana, 57
Isopterygium antarcticum, 110, 372; elegans, 57; pulchellum, 57, 334+, 372
Isothecium insidiosum, *211*; myosuroides, 57, 358+; myosuroides, distribution map for N. America, 363; stoloniferum, *358+*
isozyme banding patterns, in Cetraria arenaria, 129+

Jan Mayen (island), new bryophyte records, 332
JANSSENS, J. A. Past and Present Record of Drepanocladus crassicostatus, sp. nov. (Musci: Amblystegiaceae) and the Status of D. trichophyllus in North America, 44
Jungermannia atrovirens, 339; borealis, 334; muelleri, 339; oblongifolia, *334*; polaris, 334+; porella. *144*; pumila, 339; sphaerocarpa, 339; subelliptica, 339

KEENER, C. S. (review) E. O. Wiley. Phylogenetics. The Theory and Practice of Phylogenetic Systems, 400
KEENER, C. S. (review) V. Grant. Plant Speciation. 2nd. Edition. 176
KERSHAW, K. A. & S. WATSON. The Control of Seasonal Rates of Net Photosynthesis by Moisture, Light and Temperature in Parmelia disjuncta Erichs., 31
Kiaeria blyttii, 57; falcata, 57, 332; glacialis, 57, 334; starkei, 57, 114, 334+
Kindbergia praelonga, 57
KUMAR, S. S., see VERMA, S. K. & S. S. KUMAR
KUTA, E., see PRZYWARA, L. & E. KUTA
KUWAHARA, Y. Metzgeria **subundulata** (Aust. ex Lindb.) Kuw., stat. nov. from Texas, 276

lactate dehydrogenase, 129+
LANG, G. E., see WIEDER, R. K. & G. E. LANG
Lasallia pensylvanica, 19
LAWREY, J. D. Vulpinic and Pinastric Acids as Lichen Antiherbivore Compounds: Contrary Evidence, 365
Lecanora arctica, *17*; badia, 19; bennettii, *17*; cenisea, 19; cf. sorediza, 19; coilocarpa, 19; dispersa, 19; epibryon, 19; frustulosa, 19; intricata, 19; occidentalis, 19; polytropa, 19; proserpens, *17*; rupicola, 19; straminea, 19; superfluens, 19, 16+
Lecidea aenea, 19; aethaleoides, *18*; aglaea, 19; armeniaca, 19; assimilata, 19; atrobrunnea, 19; atromarginata, 19; auriculata, 19; berengeriana, 19; cf. lyngeana, 19; crassipes, 19; cuprea, 19; glaucophaea, 19; impavida, 19; lapicida, 19; lapicida f. ochracea, 19; marginata, 19; pantherina, 19; paupercula, 20; plana, 20; subauriculata, *19*; tessellata, 20; vernalis, 20; vorticosa, 20

Lecidella carpathica, 20; glomerulosa, 20; stigmatea, 20
Leersia contorta f. brevifolia, 336
Leiocolea collaris, 339; heterocolpos, 337+; heterocolpos var. harpanthoides, 337
Lepidoma demissa, 20
LEPP, N. W. & Y. HOCKENHULL. Growth Responses of Marchantia polymorpha Gemmalings in Relation to Concentration and Chemical Form of Applied Nickel, 342
Leptobryum pyriforme, 4, 57
Leptodictyum laxirete, 4; riparium, 4, 57; trichopodium, 57
Leptodon smithii, 110
Leptodontium viticulosoides var. panamense, *156*; L. viticulosoides var. **sulphureum**, comb. nov., *156*
Leptogium cyanescens, 119; lichenoides, 20; minutissimum, 20; saturninum, 20; tenuissium, 119
Lescuraea patens, 57, 334; radicosa, 57, 333+; radicosa var. denudata, 338
Leskea compressa, *219*
Leskeella nervosa, 57, 278; tectorum, *60*
Letharia columbiana, 368; vulpina, 365+
Leucobryum glaucum, 57
Leucodon brachypus var. andrewsianus, 57; sciuroides, 60
Leucoloma serrulatum, 378
light intensity, effects on bryophyte abundance, 1+
Limprichtia revolvens, *60*
Lobaria linita, 20; oregana, 309+
Loeskeobryum brevirostre, *60*
Loeskypnum badium, *60*; wickesii, *60*
Lopadium pezizoideum, 20
Lophozia alpestris, 115; bicrenata, 115; excisa, 335; floerkei, 339; groenlandica, 333+, kunzeana, 114; lycopodioides, 115; ventricosa, 333+, wenzelii, 114+, 338

Macromitrium cirrosum, 378; fragilicuspis, 378
Macvicaria, 144
Madotheca porella, *144*
malate dehydrogenase, 129+
Mannia fragrans, 4
MARAVOLO, N. C., F. D. BOWERS & S. FRECKMANN. The Second Annual Midwest Bryological Foray, 278
Marchantia polymorpha, 241, 333; gemmaling growth responses to nickel, 342+
MARCOTTE, F. see SHAW, J. & F. MARCOTTE
Marsupella emarginata, 241; ustulata, 115
Masonhalea richardsonii, 16+
Mazosia melanophthalma, 347+; phyllosema, 347+
Meesia muelleri, 110; triquetra, 57, 278; uliginosa, 57, 115
Megalostylium brevisetum, 124
Melanaspicilia crystallifera, *17*; dubia, 17; echtolechoides, 18; injucunda, 17; microplaca, 18; tschuctschorum, *17*
merochlorophaeic acid, 64+, 251
Meteoridium remotifolium, 379
4-*O*-methylcryptochlorophaeic acid, 64+, 251
Metzgeria bractifera, 276; furcata subsp. subundulata, 276; section Serpentina, 276; **subundulata**, stat. nov., 276; undulata, 276
Mielichhoferia elongata, 57, 274+; macrocarpa, 57, 273+, distribution map of, 274; mielichhoferi, 275; porsildii, *273*; tehamensis, 275
Missouri, new bryophyte records, 357+
Mnium ambiguum, 57; arizonicum, 114+; blytii, 114; cuspidatum, 1+; hornum, 57; lycopodioides, *60*; marginatum, 57; stellare, 57; thomsonii, 57
Moerckia blyttii, 115
Molendoa sendtneriana, 57
Monoclea forsteri, 241
Morinia saitoana, *156*; **stenotheca**, comb. nov., 156
MURRAY, B. M. (review) V. D. Aleksandrova. The Arctic and Antarctic: Their Division into Geobotanical Areas, 94
MURRAY, D. F. & B. M. MURRAY. (review) L. C. Bliss, O. W. Heal and J. J. Morre (Eds.) Tundra Ecosystems: A Comparative Analysis, 401
Mycoblastus affinis, 20; sanguinarius, 20
Myurella julacea 57, 334; sibirica, 57

Naiadita, 234
Nardia geoscyphus, 338
Neckera cladorrhizans, *219*; complanata, 57; flavescens, *229*; luridus, *232*; orthocarpus, *211*; pennata, 57, 279; sullivantii, 219
necrology, Charles Hébant, 158+; Charles M. Boardman 162
Neohyophila, 134+; lingulata, *134*; spathulifolia, *134*; sprengelii, *134*; sprengelii var. **stomatodonta**, stat. nov., 134+, 136*; sprengelii var. sprengelii, 134+, 136*
Nephroma arcticum, 20; expallidum, 20; helveticum, 119
Nevada, Porella cordaeana new to, 143+
New Zealand, Seligeria in, 106+
Newfoundland (island), checklist of the mosses of, 54+
Nicaragua, new moss records, 378+
nickel, growth responses of Marchantia polymorpha gemmalings to, 342+
norstictic acid, 17+, 254
North America, west of the hundredth meridian, the genus Porella in, 143+
Northwest Territories, Canada, lichens from the Bathurst Inlet Region, 14+

Ochrolechia androgyna, 20; frigida, 20; geminipara, 20; gonatodes, 20; grimmiae, 20; upsaliensis, 20
Odontoschisma, 150; denudatum, 279

Oediopodium griffithianum, 57
Oligotrichum hercynicum, 57
Omphalodiscus decussatus, 20; virginis, 20
Oncophorus virens, 57; wahlenbergii, 57, 339
orcinol meta-depsides, 64+
Oregon, Porella fauriei new to, 143+
Orphniospora atrata, 20
Orthodicranum flagellare, 57; montanum, 57
Orthostichidium pentagonum, 378
Orthothecium strictum, 57, 334+
Orthotrichum alpestre, 57; anomalum, 57, 279; cupulatum, 110; gymnostomum, 57; obtusifolium, 57; pallens, 57; pylaisii, 58, 336; rupestre, 114; sordidum, 58; speciosum, 58; stellatum, 58
OTTO, G. F. Tholurna dissimilis Well Established in Western North America, 263
Oxystegus tenuirostris, 337

Palamocladium nilgheriense, 110
Paludella squarrosa, 58
Papillaria nigrescens, 379
Paraleucobryum longifolium, 58
Parmelia bolliana, 119; disjuncta, 20, 246+; disjuncta, seasonal rates of photosynthesis in, 31+; elegantula, 247+; exasperatula, 20, 247+; flaventior, 246+; glabratula ssp. fuliginosa, 309; hypoleucites, 119; infumata, 20; omphalodes, 20; panniformis, 20; rudecta, 119; saxatilis, 20; septentrionalis, 20; stygia, 20; subaurifera, 119, 246+; subolivacea, 246+; sulcata, 20
Parmelina aurulenta, 119; galbina, 119
Parmeliopsis ambigua, 20
Parmotrema margaritatum, 119; michauxianum, 119
PEARD, J. L. Distribution of Corticolous Noncrustose Lichens on Trunks of Rocky Mountain Junipers in Boulder County, Colorado, 244
Pellia epiphylla, 241
Peltigera aphthosa, 20; aphthosa f. complicata, 20; canina, 20, 119; canina f. innovans, 20; canina f. sorediata, 20; canina var. rufescens, 20; collina, 20; malacea, 20; occidentalis, 14+, 20; polydactyla, 20, 40; praetextata, *20*, 40; rufescens, 41; scabrosa, 20; spuria, 119; venosa, 20
peroxidase, 129+
Pertusaria coriacea, 20; dactylina, 20; excludens, 14+, 20; monogona, 20; panyrga, 21
Petallophyllum ralfsii, 241
Phaeoceros laevis, 4; laevis subsp. carolinianus, 241
Phaeophyscia adiastola, 119; cernohorskyi, 119; ciliata, 119; erythrocardia, 119; hirsuta, 245+; hirtella, 119; orbicularis, 119; pusilloides, 119; rubropulchra, 119; sciastra, 21, 119
Phascum cuspidatum, 4, 279
Philonotis fontana, 58; marchica, 58; seriata, 338; tomentella, 334+; yezoana, 58
photosynthesis, in prairie mosses, 1+; seasonal rates in Parmelia disjuncta, 31+
phylogeny, hepatics, blepharoplast morphology in, 234+
Physcia adscendens, 21, 119, 247+; aipolia, 21, 119, 247+; americana, 119; caesia, 21, 245+; chloantha, 119; dubia, 21, 247+; grisea, 246+; millegrana, 119; orbicularis, 249+; phaea, 119; stellaris, 119, 247+
Physciopsis adglutinata, 119; syncolla, 119
Physcomitrium hookeri, 4; pyriforme, 4
Physconia detersa, 119, 247; enteroxantha, *247*; grisea, 247; muscigena, 21; pulverulenta, 247+
Pilotrichella rigida, 379
pinastric acid, 365+
Placynthium aspratile, 21
Plagiobryum zierii, 58, 334+
Plagiochila porelloides, 114
Plagiomnium ciliare, 58; cuspidatum, 58, 115; ellipticum, 58; medium, 58
Plagiopus oederiana, 58, 279
Plagiothecium cavifolium, 58, 333+; denticulatum, 58, 333+; laetum, 58, 335; latebricola, 58; platyphyllum, 60; ruthei, *60*; succulentum, 335; sylvaticum, 335
Platismatia glauca, 21
Platydictya confervoides, 279; jungermannioides, 58, 336; subtile, 58
Platygyrium repens, 58
Plaubelia tortuosa, *134*
Pleuridium ravenelii, 357; subulatum, 357; sullivantii, 357
Pleurozium schreberi 58, 334+
Pogonatum alpinum, *60*; dentatum, 58, 339; pensilvanicum, 58; urnigerum, 58, 335+
Pohlia andalusica, 58, 338; annotina, 58; bulbifera, 58; camptotrachela, 60; cruda, 58, 114, 333+; cucullata, 337; drummondii, 58, 334; lescuriana, 58; nutans, 58, obtusifolia, 58, 114+, 337; proligera, 58, 338; pulchella, *60*; rothii, *60*; sphagnicola, 58; wahlenbergii, 58, 338
Polytricastrum alpinum, 333; alpinum, 58, 335; formosum, 58; longisetum, 58, 115; lyallii, 114; pallidisetum, 58; sexangulare, 58, 333+
Polytrichum affine, *60*; alpestre, 284; alpinum, 335; capillare, 339; commune, 58, 280+; formosum, *60*; hyperboreum, 335; juniperinum, 2, 58, 281+, 338; longisetum, *60*; ohioense, 60, 281; piliferum, 58, 335; strictum, 58, 335
population dynamics and production, in a population of Lobaria oregana, 309+
Porella bolanderi, 145+; bolanderi, map of N. American distribution, 147; caespitans, 152; cordaeana, 145+; cordaeana, map of N. American distribution, 147; fauriei, 145+; navicularis, 145+; navicularis, map of N. American distribution, 151; pinnata, 144; pla-

typhylla, 145+; platyphylloidea, 145+; platyphylloidea, map of western N. American distribution, 147; roellii, 145+; roellii f. **schofieldiana**, fo. nov., 153; roellii fo. **crispata**, fo. nov., 153; roellii, map of N. American distribution, 151; setigera, *152*; thuja, 150; ulophylla, 146+; vernicosa, 154; in North America west of the hundredth meridian, 143+; **Navicularis**, subsect. nov., 145+; subgenus Macvicaria, 144; subgenus Porella, 144; subgenus Protoporella, 144; subsection Platyphyllae, 145+
Porina epiphylla, 347+; impressa, 348+; obducta, 348+; semicarpi, 348+
porphyrilic acid, 21
Pottia davilliana, 4; latifolia, *60*
Pottiaceae, 156, 134+, 271+
prairie bryophytes, 1+
PRENDERGAST, H. D. V. Three Mosses New to Upper Volta, 270
protocetraric acid, 66, 254
protolichesterinic acid, 64+
PRZYWARA, L. & E. KUTA. An Acetic-Haematoxylin Method for Cytological Investigations of Bryophyta, 141
Pseudephebe minuscula, 21; pubescens, 21
Pseudobryum cinclidioides, 58
Pseudoleskea breidleri, 338; patens, *60*
Pseudoleskeella catenulata, 58, 278; papillosa, 61; tectorum, 58; williamsii, 61
Pseudoparmelia caperata, 119
Pseudoscleropodium purum, 58
Pseudostereodon procerrimum, *61*
Psora decipiens, 21; rubiformis, 21
Psoroma hypnorum, 21
psoromic acid, 254
Pterigynandrum filiforme, 58
Pteryobryopsis mexicana, 379
Ptilium crista-castrensis, 58
Pylaisiella intricata, 58; polyantha, 58, 278; selwynii, 58
Pyxine sorediata, 119

Racomitrium aciculare, 58; canescens, 58, 124, 336+; canescens var. ericoides, 127; ericoides, 333+; fasciculare, 58, 333; heterostichum, 58, 333+; lanuginosum, 58, 333+; microcarpon, 58, 338+; patens, 127; sudeticum, 58, 338+
Racopilum tomentosum, 378
Radula complanata, 114
Ramalina americana, 119
rangiformic acid, 64+
REDFEARN, P. L., JR. Bryophytes of the Interior Highlands of North America XVIII. Additions to the Flora, 357
REDFEARN, P. L. (review) E. V. Watson. British Mosses and Liverworts. 3rd Edition, 176
REESE, W. D. American Calymperes and Syrrhopodon: Identification Key and Summary of Recent Nomenclatural Changes, 23
REESE, W. D. Syrrhopodon brasiliensis, a Previously Undescribed Species from Southern Brazil, 354
RENLIANG, H. A Revision of the Chinese Species of Entodon (Musci, Entodontaceae), 193
Retidens, *194*
Rhabdoweisia crispata, 58
Rhacomitrium aciculare, 357; heterostichum var. occidentale, 115; sudeticum f. alpinum, 114
Rhizocarpon copelandii, 21; disporum, 21; eupetraeum, 21; geographicum, 21; grande, 21; inarense, 21; superficiale, 21, 42
Rhizogonium spiniforme, 378
Rhizomnium appalachianum, 58; magnifolium, 58; pseudopunctatum subsp. chlorophyllosum, 58; punctatum, 58
Rhizoplaca melanopthalma, 21
RHOADES, F. M. Distribution of Thalli in a Population of the Epiphytic Lichen Lobaria oregona and a Model of Population Dynamics and Production, 309
Rhodobryum ontariense, *61*; roseum, 278; spathulatum, 58
Rhynostegiella compacta, *61*
Rhytidiadelphis triquetrus, 58; loreus, 58; squarrosus, 58, 261, 336; subpinnatus, 58; triquetrus, 278
Rhytidium rugosum, 58
Riccia fluitans, 4
Ricciocarpus natans, 4
Riella americana, 241, blepharoplast structure in, 97+
Rinodina cacuminum, 21; milvina, 21; mucronulata, 21; oreina, 33+; septentrionalis, 21; turfacea, 21
Roellia roellia, 2
ROGERS, W., see CONRAN, J. G. & R. W. ROGERS
RUDOLPH, E. D. (review) J. Poelt & A. Vezda. Bestimmungsschlüssel europäischer Flechten. Ergänzungshelt II., 95

Saelania glaucescens, 58, 335
Sakuraia, *194*; conchophyllus, *223*; macrospora, *223*
salazinic acid, 254
Sanionia uncinata, *61*
Scapania curta, 333; hyperborea, 338; irrigua, 338; obcordata, 338; umbrosa, 114
Schistidium agassizii, 58; alpicola, 61; apocarpum, 58, 338; confertum, 338; gracile, 58; maritimum, 58; maritmum var. maritmum, 335; rivulare, 58; rivulare subsp. latifolium, 338; strictum, 338; trichodon, 58
Schistostega scorpioides, 58; turgescens, 58
Schlotheimia rugifolia, 378
SCHUTTE, J. A. Distribution Patterns of Foliose and Fruticose Lichens of Eastern Iowa, 117
Scleropodium obtusifolium, 114

Scorpidium vernicosum, *61*
SCOTTER, G. W., see THOMSON, J. W. & G. W. SCOTTER
sekikaic acid, 64+
Seligeria, in New Zealand, 106+; brevifolia, 58, 109; calcarea, 58; campylopoda, 58, 111, 279; cardotii, 372, 106+, 108*; careyana, 109; diminuta, 106+, 112*; diversifolia, 111; donniana, 59; pusilla, 106+; recurvata, 59, 106+; tristichoides, 59
Sematophyllum marylandicum, 59
SHAW, J. & F. MARCOTTE. Bryum warneum in North America, 266
Siphula ceratites, 21
SNIDER, J. A. (review) R. E. Andrus. Sphagnaceae (Peat Moss Family) of New York State., 191
SNIDER, J. A., see ALLEN, B. H. & J. A. SNIDER
SOLLMAN, P. Notes on Pottiaceous Mosses I., 271
Solorina crocea, 21; saccata, 21
South Dakota, Porella platyphylla new to, 143+
Southbya tophacea, 241
Sphaerocarpales, comparative spermatogenesis in, 97
Sphaerocarpos donnellii, 241; donnellii, blepharoplast structure in, 97
Sphaerophorus fragilis, 21; globosus, 21
Sphagnum angermannicum, 59; angustifolium, 59, 281; annulatum, 261; annulatum var. porosum, 59; balticum, 59, 284; capillaceum, *61*; capillifolium, 59; **carolinianum**, 257+, 258*, 259*, 261*; centrale, 59; compactum, 59; contortum, 59, 261; **crispum**, 257+, 258*, 259*, 261*; cuspidatum, 59, 284; fallax, 59, 281+; fimbriatum, 59; fitzgeraldii, 260; flavicomans, 59; flexuosum, *61*, 281; fuscum, 59, 284, 357; girgensohnii, 59, 281; henryense, 357; imbricatum, 59, 281; lescurii, 257+; lindbergii, 59; macrophyllum, 260; macrophyllum var. burinense, 59; magellanicum, 59, 280+; majus, 59; mendocinum, 261; molle, 59; nemoreum, 115, 261; palustre, 59; papillosum, 59, 281+; platyphyllum, 59; pulchrum, 59, 260, 284; pylaesii, 59; quinquefarium, 59; recurvum, 59, 280+; riparium, 59; rubellum, 59, 284; russowii, 59, 261, 357; squarrosum, 59; strictum, 59; subfulvum, 59; subnitens, 59; subsecundum, 59, 257+, 259*, 281; tenellum, 59; tenerum, 59, 260, 357; teres, 59; torreyanum, 59, 260; warnstorfii, 59; wulfianum, 59
Splachnum ampullaceum, 59; ovatum, *61*; sphaericum, 59; vasculosum, 59
Sporastatia testudinea, 21
Sporopodium phyllocharis var. flavescens, 348+; xantholeucum, 347+
squamatic acid, 254
Squamulidium nigrescens, 379
STARK, L. R. (review) A. J. E. Smith, Editor. Bryophyte Ecology, 306
Stegonia latifolia, 59
Stereocaulon alpinum, 21; arenarium, 21; leprocephalum, 21; paschale, 40; rivulorum, 21; symphycheilum, 21; tomentosum, 21
Stereodon caligino sus, *213*; luridus, *232*; prorepens, *225*; thomsonii, *225*
Stereophyllum fulvum, 127; tavoyense, 127
Sticta arctica, 21, 16+
stictic acid, 254
Stokesiella praelonga, *61*; praelonga var. praelonga, 115
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature—97–100: 74, 163, 287, 380
Streblotrichum convolutum, *61*
Strigula complanata, 347+; elegans, 348+; subtilissima, 347+
Synodontella, *194*
Syrrhopodon annotinus, 25+; **brasiliensis**, sp. nov., 354+, 355*; circinatus, 28; cryptocarpos, 24+, 27; cymbifolius, 26; elongatus, 25; elongatus var. elongatus, 25; elongatus var. glaziovii, 25; erubescens, 25; fimbriatus, 25; flexifolius, *30*; gardneri, 124; gaudichaudii, 26, 355; helicophyllus, 25; hornschuchii, 28; incompletus, 24+; incompletus var. berteroanus, 28; incompletus var. elatus, 28; incompletus var. incompletus, 28; incompletus var. lanceolatus, 24+; incompletus var. luridus, 28; incompletus var. perangustifolius, 24+; isthmi, 26; leprieurii, 25; ligulatus, 26; lycopodioides, 26; parasiticus var. disciformis, 26; parasiticus var. flexifolius, 26; parasiticus var. parasiticus, 26; prolifer, 356; prolifer var. acanthoneuros, 27; prolifer var. cincinnatus 25+; prolifer var. papillosus, 27; prolifer var. prolifer, 27; prolifer var. tenuifolius, 27; quintasii, 24+; ramicola, *30*; rigescens, 25; rigidus, 24+; rivularis, *30*; rubiginosus, *30*; rupestris, 27; simmondsii, 25; steyermarkii, 27; texanus, 24; theriotii, 24+, ulei, 25+, 356; xanthophyllus, 27

Tayloria lingulata, 59; serrata, 59; splachnoides, 59
Tetradontium brownianum, 59; repandum, 59
Tetraphis geniculata, 59; pellucida, 59
Tetraplodon angustatus, 59; blyttii, 335; bryoides, 335; mnioides, 59; pallidus, 59
Tetradontium repandum, 335
Thamnolia subuliformis, 21
thamnolic acid, 254
Tholurna dissimilis, 263+
THOMSON, J. W. & G. W. SCOTTER. Lichens from Bathurst Inlet Region, Northwest Territories, Canada, 14
Thuidium delicatulum, 59; philbertii, 59; recognitum, 59; tamariscinum, 59
Timmia austriaca, 59, 339, 370; bavarica, 370; comata, *370*; megalopolitana, 370; norvegica, 370+; norvegica var. excurrens, 59, 370+; norvegica var. norvegica, 59, 370+; siberica, 370

Tomentypnum falcifolium, 59; nitens, 59, 338
Toninia caeruleonigricans, 21; squalida, 21
Tortella arctica, 59; fragilis, 59, 335, 372; tortuosa, 59, 278
Tortula berteroana, *134*; donellii, *134*; mucronifolia, 59, 335; muralis, 59, 110; norvegica, 59, 114; phaea, 110; ruralis, 278, 336; ruralis var. crinita, 59; ruralis var. ruralis, 59; sprengelii, *134*; vinealis, 272
Trematodon ambiguus, 59; montanus, 59
Tremolechia atrata, 21
Tricharia albostrigosa, 347+
Trichodon cylindricus, 59
Trichostomum berteroana, *134*; berteroi, *134*; crispulum, 59; sprengelii, *134*; sulphureum var. panamense, *156*; sulphureum var. sulphureum, *156*; tenuirostre, 59, 337, 378
Tridontium tasmanicum, 110
Tritomaria polita ssp. acuta, gemmae of, 139+; polita ssp. polita, gemmae of, 139+, 140*; polita ssp. polymorpha, gemmae of, 140*; quinquedentata, 335; scitula, 139+, 338

Ulota coarctata, 59; crispa, 8, 59; curvifolia, 59; drummondii, 59; hutchinsiae, 59; phyllantha, 59
Umbilicaria arctica, 33+; cylindrica, 21; deusta, 21, 41; havaasii, 21; hyperborea, 21; mammulata, 41; papulosa, 41; proboscidea, 21; torrefacta, 21; vellea, 21, 41
Upper Volta, mosses new to, 270
Usnea arizonica, 255; florida, 254; hirta, 119, 247; **lecanorica**, sp. nov., 254+, 255*; subfloridana, 309; trichodea, 254
usnic acid, 254, 368

VAN DER LINDEN, J. O. & D. R. FARRAR. An Ecological Study of the Bryophytes of a Natural Prairie in Northwestern Iowa, 1
VERMA, S. K. & S. S. KUMAR. Cytological Observations on Some West Himalayan Mosses—VIII, 121
VITT, D. H. & J. K. BARTLETT. The Genus Seligeria in New Zealand, 106
vulpinic acid, 365+

Warnstorfia exannulata, *61*; procera, *61*
Washington, state, western, the Cladonia chlorophaea-C. fimbriata complex in, 64+
WATSON, S., see KERSHAW, K. A. & S. WATSON
Weisiopsis oblonga, 134+; spathulifolius, *134*; stomatodonta, *138*
Weissia berteroana, *134*; controversa, 1+, 59, 279; planifolia, 156; rutilans, 156; sprengelii, *134*; **subangustifolia**, comb. nov., 156
WIEDER, R. K. & G. E. LANG. Net Primary Production of the Dominant Bryophytes in a Sphagnum-Dominated Wetland in West Virginia, 280
Wilkiea macrophylla, lichen succession on, 347+
winter hardening, in lichens, 31+

Xanthoparmelia centrifuga, 21; incurva, 21; separata, 21; taractica, 22
Xanthoria candelaria, 22, 119; elegans, 22; fallax, 119, 245+; polycarpa, 119, 247; sorediata, 22

ZANDER, R. H. Nomenclatural Changes in Hyophila, Leptodontium and Morinia (Pottiaceae), 156
ZANDER, R. H. A Reevaluation of Neohyophila Crum (Pottiaceae), 134
Zelometeorium allionii, 378+
Zygodon conoideus, 59; viridissimus, 59; vulgaris, *61*